

D. Steyer, P. Tristram, C. Clayeux, F. Heitz and B. Laugel

# Yeast Strains and Hop Varieties Synergy on Beer Volatile Compounds

The volatile compounds produced in beers are complex and influenced by numerous factors like barley, brewing and fermentation process, hop variety and yeast strain. In this study, hop variety and yeast strain have been chosen to evaluate their influences on the volatile compound profile in beer. Malt wort was hopped with 4 different varieties of French aromatic hops and fermented with 3 different yeast strains. Volatile compounds were analysed by Stir Bar Sorptive Extraction-Gas-Chromatography (SBSE-GC-MS). Statistical analysis (ANOVA) were performed to test the influence of yeast strain and hop variety on the concentration of each of the measured compounds. Among the 39 volatile compounds detected, 9 of them were influenced only by hop variety, 2 by yeast only and the 28 remaining by both hop and yeast together. As expected, terpenes concentration in beer is directly linked to the choice of hop variety used, but it appears the yeast strains can also influence that parameter. Surprisingly, hops variety influenced the final concentration of esters, highlighting that an interaction between hops compounds and yeast metabolism remains to be investigated.

Descriptors: hop varieties, yeast strains, volatile compounds

## 1 Introduction

The final taste of beer is a sum of the various aromas provided by raw materials during the different steps of the beer production process. The choice of yeast strain and hop variety are often considered as determinants of the beers aromatic profile.

Among all the volatile compounds involved in the quality of beer, yeast is involved in the production of some important flavour-active compounds like isoamyl acetate, 2-phenylethanol, 4-vinylguaïacol or ethyl hexanoate as reviewed by *Carrau et al.* [4]. The final concentration of these compounds is yeast-strain dependant [20, 39], diversity of brewer's yeast helps brewers to drive the aroma of beer.

In addition to yeast strain, hop variety is the other factor that brewers can also choose to strongly affect the aroma of beer [19]. Most of the flavour active-compounds brought by aromatic hops are monoterpenes, sesquiterpenes, esters and alcohols [29, 40]. Currently, only monoterpenes, especially geraniol, nerol and citronellol, are known to be influenced by yeast metabolism during fermentation [21], while sesquiterpenols, esters and alcohols are only suspected to be bioconverted by yeast.

<https://doi.org/10.23763/BrSc17-13Steyer>

### Authors

Damien Steyer, Pauline Tristram, Céline Clayeux, Twistaroma, Colmar, France; Francis Heitz, Bernadette Laugel, Comptoir Agricole/Cophoudal/AGPH, Hochfelden, France; corresponding author: damien.steyer@twistaroma.fr

To investigate the influence of hop varieties and yeast strains on the volatile content of beer, four hop varieties from Cophoudal/Comptoir Agricole and two different brewer's yeast strains from the CLIB (Institut Pasteur/INRA collection) and one from the University of California, Davis (UCD) have been used to produce beer in triplicate. Following this, the beers were analysed by SBSE-GC-MS to determine their final volatile composition and to evaluate the influence of the hop variety and the yeast strain, but also the combination of both.

## 2 Materials and methods

### 2.1 Hop assays

Four French aromatic hop varieties harvested in 2013 were used to produce hopped wort, namely Strisselspalt, Aramis, Bouclier and Triskel (Cophoudal/Comptoir Agricole Hochfelden, France). Table 1 shows the linalool and  $\alpha$ -acids composition of each hop variety.

Twenty litres of wort were brewed for each hop variety with malt extract (Bières du monde, Lille, France) and distilled water, heated to 100 °C, to obtain wort at 13 °Plato with a Braumeister 50 L (Speidel, Offerdingen, Germany). Hop (2,1 g/L) was added in a tea bag (10 min, 100 °C). After tea bag removal, wort was quickly cooled down to 20 °C and let at room temperature during the preparation of the fermentation.

Table 1 Hop varieties composition

Compounds	Hop varieties			
	Strissel-spalt	Aramis	Bouclier	Triskel
Linalool	7–8 mg/g	14 mg/g	6 mg/g	10 mg/g
$\alpha$ -acids	3.4%	8.1%	9.5%	8.7%

**Table 2** Results of the two ways Anova for the two factors Hop and Yeast and their possible interactions

Compounds	Hop	Yeast	Yeast/hop
<b>Acetate Esters</b>			
Hexyl acetate	0.000	0.000	0.014
2-Phenethyl acetate	0.026	0.000	0.009
Isoamyl acetate	0.020	0.000	0.005
<b>Medium Chain fatty acids</b>			
Octanoic acid	0.000	0.000	0.038
Decanoic acid	0.000	0.000	0.000
Dodecanoic acid	0.000	0.000	0.000
<b>Alcohols</b>			
2-Hexanol	0.000	0.000	0.000
2-undecanol	0.000	0.000	0.000
3-methyl-2-Buten-1-ol	0.000	0.055	0.168
2-Nonanol	0.000	0.880	1.000
<b>Esters</b>			
2-Phenylethyl butanoate	0.000	0.000	0.000
2-Phenethyl hexanoate	0.050	0.000	0.016
Isoamyl propanoate	0.000	0.001	0.000
Methyl 4-Methyl-2-hexenoate	0.000	0.031	0.000
Methyl geranate	0.000	0.043	0.739
isobutyl butanoate	0.000	0.643	0.954
Ethyl hexanoate	0.000	0.000	0.000
Ethyl octanoate	0.000	0.000	0.000
Ethyl decanoate	0.000	0.000	0.000
Ethyl dodecanoate	0.000	0.000	0.000
<b>Higher Alcohols</b>			
2-Phenyl Ethanol	0.000	0.000	0.000
Isoamyl alcohol	0.489	0.000	0.575
<b>Monoterpenes</b>			
2,7-Dimethyl-2,7-octadiene	0.000	0.060	0.417
Citronellol	0.000	0.000	0.001
Nerol	0.000	0.000	0.000
Linalool	0.000	0.086	0.021
Geraniol	0.000	0.000	0.000
<b>Sesquiterpenols</b>			
7-epi-amiteol	0.000	0.000	0.000
a/t-Cadinol	0.000	0.000	0.000
Hop ether	0.000	0.090	0.017
Juniper Camphor	0.000	0.447	0.438
$\beta$ -Eudesmol	0.000	0.008	0.049
$\alpha$ -Eudesmol	0.000	0.010	0.055
Caryophyllene oxide	0.000	0.837	0.601
<b>Terpenyl acetates</b>			
Citronellol acetate	0.000	0.040	0.050
Geranyl acetate	0.000	0.575	0.217
Neryl acetate	0.000	0.000	0.000
<b>Phenols</b>			
4-Vinylguaiacol	0.000	0.000	0.000
Styrene	0.831	0.000	0.121
<b>Other</b>			
2,3-dihydro-benzofuran	0.041	0.000	0.014

## 2.2 Yeast assays

Three yeast strains were chosen to ferment the wort, one *S. cerevisiae* ale strain UCD915, and two lagers CLIB267 and CLIB279, kindly provided by Jean-Luc Legras from the INRA Colmar collection.

Yeast pre-cultures were performed on YPD medium (Yeast extract 10 g/L, peptone 20 g/L, glucose 20 g/L) at 28 °C for 24 h in 125 mL flasks under shaking. Beer wort was inoculated at initial  $OD_{600nm} = 0.1$ . Fermentation steps were performed at 15 °C in 1 L autoclaved vessels equipped with airlocks to maintain anaerobiosis, with constant stirring (250 rpm). Vessels were weighed twice daily for fermentation monitoring, and fermentation was stopped when the daily loss reached less than 1 % of the expected total loss. Each of the 4 different worts were fermented with each yeast strain in triplicate, to a total of 36 different beer combinations.

## 2.3 Volatile analysis

Analysis of the volatiles compounds of the 36 beers was performed using the Stir Bar Sorptive Extraction method-Liquid Desorption Gas Chromatography [6] (SBSE-LD-GC-MS). Each beer was analysed in duplicate. All reagents used were analytical grade. Stir Bars (length = 20 mm) were coated with 47  $\mu$ L of polydimethylsiloxane (Twister; Gerstel, Mülheim a. d. Ruhr, Germany). The GC-MS analyses were performed with an Agilent 6890N gas chromatograph equipped with an Agilent 7683 automatic liquid sampler coupled to an Agilent 5975B inert Mass Spectrometer Detector (Agilent Technologies). The gas chromatograph was fitted with a DB-Wax capillary column (60 m  $\times$  0.32 mm i.d.  $\times$  0.50  $\mu$ m film thickness, J&W Scientific) with helium carrier gas (1 mL/min, constant flow). Instrument control and data processing were performed with Agilent MSD ChemStation software (G1701DA, Rev D.03.00). The mass spectra were compared with the Wiley's library reference spectral bank, Retention Index (RI) and standard when available. All compounds were semi-quantified using the ratio of their Total Ion Current peak to that of the 3-octanol (final concentration of 84  $\mu$ g/L).

## 2.4 Statistical analyses

Statistical analyses were performed on Minitab 17.0 using Anova.

## 3 Results and discussion

The 36 beers produced have been analysed to investigate the influence of hop varieties, yeast strains and their potential interactions on 39 volatile compounds (Table 2).

Anova statistical analysis indicated that among the identified compounds, 9 were influenced by hop only, 2 by yeast only the 28 remaining were both influenced by hop and yeast. Most of the compounds influenced by the synergistic effect of yeast and hop (interaction yeast/hop in Table 2) are also influenced by hop and yeast separately. This is why these interaction are not discussed in this study.

A schematic representation of the conclusion of this study is presented in figure 1.

### 3.1 Impact of hop variety on beer volatile profile

The 9 compounds influenced by the hop variety only include three monoterpenes: linalool (1), geranyl acetate (2), 2,7 dimethyl 2,6 octadiene (3); three sesquiterpenols: hop ether (4), caryophyllenoxyde (5) and juniper camphor (6); one ester: isobutyl isobutanoate (7) and two alcohols: 2-nonanol (8) and 3-methyl-2-Buten-1-ol (9). Among all of the compounds influenced by hop variety, linalool (1) is a key contributor to hop aroma in beer [16]. Its perception threshold in beer is relatively low (27–100 µg/L [7]) and its descriptors are floral, citrus and flower. Previous studies showed that yeast was able to metabolize geraniol, citronellol and nerol but not linalool [22]. As expected, in our conditions, the concentration of this compound in beer depended only on the hop variety used. Among the four hops tested, the highest concentration of linalool was obtained with Aramis (≈ 500 µg/L rel 3-octanol, Fig. 2).

Geranyl acetate (2) is described as floral and reported to be involved in the hoppy aroma. However, as reported by *Lermusieau et al.* [26] its impact on beer aroma is still unclear. Only Aramis and Triskel hops bring geranyl acetate to beer whatever the yeast strain used (Fig. 1). This compound could be the result of the esterification of geraniol to geranyl acetate by yeast during fermentation [41] but could also come directly from hop [28, 37, 38]. Our conditions are not adapted for origin determination of this compound and need further investigation that are not the scope of this work.

Bouclier and Aramis show the highest concentration of 2,7 dimethyl 2,6 octadiene (3). This compound has already been reported in beer [1, 46] and in




Fig. 1 Schematic representation of the interaction between hop and yeast volatile compounds during the fermentation of beer (green arrows: yeast enzymatic reactions)




Fig. 2 Graphical representation of the concentration of linalool and geranyl acetate (µg/L rel 3-octanol) in beers produced with Aramis, Bouclier, Strisselspalt and Triskel as aromatic hop and fermented with yeast CLIB267, CLIB279 and UCD915




Fig. 3 Graphical representation of the concentration of Isoamyl alcohol and Styrene (µg/L rel 3-octanol) in beers produced with Aramis, Bouclier, Strisselspalt and Triskel as aromatic hop and fermented with yeast CLIB267, CLIB279 and UCD915

hop [27], but to our knowledge, there is no data about its influence on the aroma of beer.

Aramis appears to be the highest producer of hop ether (4) in our conditions. This aroma-active compound founds in hop oil in previous study [44]. Its odour impact on beer has been evaluated by Lam et al. [25] who determined that this compound was not a major contributor to hop aroma. Its influence on beer remains to be elucidated.

Caryophyllen oxide (5) brought by hop [28, 45] was known to contribute to the spicy aroma of beer [11] and was involved in the “Noble” character of hop varieties. This compound can be the result from the oxidation or hydrolysis of β-caryophyllen during the boiling step [15]. Aramis and Bouclier bring the highest concentration of this compound in beer in this study (Table 3 and Table 4).

Juniper camphor (6) has been previously reported in beer produced with Aramis and Strisselspalt [40] and was reported to be present in hop oil [2,9]. Its influence in beer aroma is not yet defined. This compound is only detected in beer hopped with Aramis (Table 3 and Table 4).

Isobutyl isobutanoate (7) has been reported in hop oil [12,31] and beer [2]. It displays a relatively low threshold perception at 36

µg/L [3] and with a positive description for the aroma of beer, ie pineapple. Aramis is the highest producer of this compound during our experiments.

2-nonanol (8) and 3-methyl-2-buten-1-ol (9) are 2-alkanols which have been reported in hop oil [10,37] and beer [2]. According to Hashimoto [17,18], they are the results of degradation of iso-α-acids of hop during the oxidation of beer. This reaction leads also to the formation of 2-alkanone which could be further reduced to 2-alkanols by yeast [47]. Their respective influences on the final aroma of beer are not known. Aramis is the highest producer of these compounds even if its α-acids is not the highest.

### 3.2 Impact of yeast strain on beer volatile profile

The two volatile compounds only influenced by yeast strains are isoamyl alcohol (10) and styrene (11) (Table 2). Isoamyl alcohol (10) concentration is statistically equivalent in beer whatever the hop variety chosen but its concentration is different depending on the yeast used in fermentation. As a by-product of the metabolism of yeast during fermentation (Fig.1) this compound is known to be dependent on the yeast strain [2,20,39]. Lermusieau et al have shown that unhopped and hopped beer (with Saaz and Challenger) have the same factor of dilution (linked to the concentration of the compounds) for isoamyl alcohol in GC-O experiment [26].

**Table 3** Concentration ( $\mu\text{g/L}$  relative to 3-octanol) of volatile compounds in beer hopped with Strisselspalt and Bouclier and fermented with CLIB 267, CLIB279 and UCD915


Hop varieties	Yeast strains	Strisselspalt						Bouclier					
		CLIB267		CLIB279		UCD915		CLIB267		CLIB279		UCD915	
Compounds	N°	Mean	Err	Mean	Err	Mean	Err	Mean	Err	Mean	Err	Mean	Err
<b>Acetate Esters</b>													
Hexyl acetate <sup>1,2</sup>	32	2.4	29%	3.5	25%	8.4	29%	12.7	5%	19.2	19%	15.5	14%
2-Phenethyl acetate <sup>1,2</sup>	33	784.8	9%	744.9	26%	2115.4	38%	938.4	6%	1248.6	5%	1624.0	4%
Isoamyl acetate <sup>1,2</sup>	31	194.3	11%	605.3	30%	1810.9	44%	699.3	10%	3150.4	9%	2901.4	5%
<b>Medium Chain fatty acids</b>													
Octanoic acid <sup>1,2</sup>	36	205.3	18%	362.6	35%	1169.7	48%	807.7	40%	1811.4	17%	2066.5	14%
Decanoic acid <sup>1,2</sup>	37	1258.9	9%	1411.1	34%	3713.3	38%	1559.9	4%	3141.5	4%	8366.5	3%
Dodecanoic acid <sup>1,2</sup>	38	0.0		53.4	183%	4.9	67%	252.0	162%	2657.4	6%	1284.3	10%
<b>Alcohols</b>													
2-Hexanol <sup>1,2</sup>	22	0.0		0.0		34.0	70%	16.0	4%	4.7	173%	0.0	
2-undecanol <sup>1,2</sup>	21	0.0		0.0		0.0		19.8	3%	13.9	13%	8.5	8%
3-methyl-2-Buten-1-ol <sup>1</sup>	9	0.0		0.0		0.0		3.3	87%	0.0		4.6	18%
2-Nonanol <sup>1</sup>	8	0.0		0.0		0.0		7.7	87%	3.3	173%	7.2	87%
<b>Esters</b>													
2-Phenylethyl butanoate <sup>1,2</sup>	34	16.0	35%	3.8	13%	6.5	40%	98.7	6%	23.6	3%	16.4	9%
2-Phenethyl hexanoate <sup>1,2</sup>	35	837.7	21%	119.4	12%	0.0		1434.8	9%	75.7	13%	41.6	9%
Isoamyl propanoate <sup>1,2</sup>	19	0.9	35%	1.5	14%	0.0		4.5	87%	5.2	95%	5.6	4%
Methyl 4-Methyl-2-hexenoate <sup>1,2</sup>	20	5.5	16%	6.3	5%	9.0	18%	91.0	7%	95.7	7%	78.9	5%
Methyl geranate <sup>1,2</sup>	21	0.0		0.0		0.0		13.9	15%	16.2	19%	14.4	11%
isobutyl butanoate <sup>1</sup>	7	0.0		0.0		0.0		3.8	89%	4.7	15%	4.6	15%
Ethyl hexanoate <sup>1,2</sup>	27	119.8	13%	99.6	9%	697.8	64%	603.4	5%	463.0	7%	1060.8	6%
Ethyl octanoate <sup>1,2</sup>	28	112.1	15%	196.1	21%	549.4	46%	570.7	2%	941.0	7%	856.7	5%
Ethyl decanoate <sup>1,2</sup>	29	31.0	35%	38.1	37%	157.3	51%	146.1	3%	316.2	8%	455.6	6%
Ethyl dodecanoate <sup>1,2</sup>	30	3.4	29%	9.6	69%	20.8	43%	39.6	10%	140.2	10%	82.0	7%
<b>Higher Alcohols</b>													
2-Phenyl Ethanol <sup>1,2</sup>	39	254.4	19%	207.8	41%	491.4	42%	939.1	5%	863.4	18%	730.3	20%
Isoamyl alcohol <sup>2</sup>	10	547.3	6%	868.9	18%	1735.9	37%	540.2	3%	1085.0	7%	1647.8	9%
<b>Monoterpenes</b>													
2,7-Dimethyl-2,7-octadiene <sup>1</sup>	3	6.1	87%	13.0	36%	0.0		84.0	49%	42.5	8%	38.1	28%
Citronellol <sup>1,2</sup>	14	0.0		0.0		0.0		23.9	8%	18.8	6%	11.7	173%
Nerol <sup>1,2</sup>	13	0.0		0.0		0.0		0.0		0.0		1.2	173%
Linalool <sup>1</sup>	1	39.9	4%	43.4	12%	43.6	7%	154.3	3%	147.1	3%	150.7	4%
Geraniol <sup>1,2</sup>	12	4.8	21%	4.2	14%	5.3	7%	12.0	1%	7.7	4%	11.5	14%
<b>Sesquiterpenols</b>													
7-epi-amiteol <sup>1,2</sup>	17	0.0		0.0		0.0		0.0		0.0		0.0	
a/t-Cadinol <sup>1,2</sup>	18	0.0		0.0		0.0		0.0		28.4	16%	0.0	
Hop ether <sup>1,2</sup>	4	0.0		0.0		0.0		4.6	173%	13.8	8%	9.1	87%
Juniper Camphor <sup>1</sup>	6	0.0		0.0		0.0		0.0		0.0		0.0	
$\beta$ -Eudesmol <sup>1,2</sup>	15	38.5	11%	32.6	22%	0.0		9.9	173%	0.0		0.0	
$\alpha$ -Eudesmol <sup>1,2</sup>	16	35.7	5%	34.3	24%	0.0		0.0		0.0		0.0	
Caryophyllene oxide <sup>1</sup>	5	19.3	3%	21.0	20%	18.6	26%	69.5	19%	63.7	41%	54.4	15%
<b>Terpenyl acetates</b>													
Citronellol acetate <sup>1,2</sup>	24	0.0		0.0		0.0		15.7	9%	15.1	10%	14.5	16%
Geranyl acetate <sup>1</sup>	2	0.0		0.0		0.0		17.2	14%	15.5	3%	10.4	5%
Neryl acetate <sup>1,2</sup>	23	0.0		0.0		0.0		10.3	5%	17.22	15%	15.46	3%
<b>Phenols</b>													
4-Vinylguaicol <sup>1,2</sup>	25	19.6	9%	15.2	13%	573.0	67%	17.9	14%	15.1	9%	411.1	10%
Styrene <sup>2</sup>	11	0.0		0.0		88.9	70%	4.3	173%	6.0	104%	76.0	13%
<b>Other</b>													
2,3-dihydro-benzofuran <sup>1,2</sup>	26	0.0		0.0		88.8	86%	0.0		0.0		19.8	87%

<sup>1</sup> compounds influenced by hop variety (p-value < 0.05); <sup>2</sup> compounds influenced by yeast (p-value < 0.05)

**Table 4** Concentration (µg/L relative to 3-octanol) of volatile compounds in beer hopped with Triskel and Aramis and fermented with CLIB267, CLIB279 and UCD915

Hop varieties	Yeast strains	Triskel						Aramis					
		CLIB267		CLIB279		UCD915		CLIB267		CLIB279		UCD915	
Compounds	N°	Mean	Err	Mean	Err	Mean	Err	Mean	Err	Mean	Err	Mean	Err
<b>Acetate Esters</b>													
Hexyl acetate <sup>1,2</sup>	32	16.0	11%	19.9	27%	30.4	5%	16.1	9%	26.2	19%	34.3	28%
2-Phenethyl acetate <sup>1,2</sup>	33	1221.1	14%	953.2	26%	2291.9	19%	988.6	17%	891.8	20%	1839.5	17%
Isoamyl acetate <sup>1,2</sup>	31	1132.5	13%	2790.3	28%	5920.2	9%	829.4	13%	2860.3	13%	5766.6	5%
<b>Medium Chain fatty acids</b>													
Octanoic acid <sup>1,2</sup>	36	1623.0	16%	2247.6	19%	4124.5	2%	1244.8	3%	2088.0	9%	3997.6	7%
Decanoic acid <sup>1,2</sup>	37	1670.5	26%	1892.2	36%	8788.1	11%	638.8	3%	2409.0	17%	9507.9	17%
Dodecanoic acid <sup>1,2</sup>	38	0	0	735.8	82%	2455.9	8%	109.0	39%	1460.0	51%	2333.1	8%
<b>Alcohols</b>													
2-Hexanol <sup>1,2</sup>	22	0.0		0.0		1.3	173%	9.5	3%	10.3	15%	11.2	18%
2-undecanol <sup>1,2</sup>	21	16.5	21%	10.8	8%	13.9	19%	67.4	5%	43.6	23%	63.1	35%
3-methyl-2-Buten-1-ol <sup>1</sup>	9	7.1	46%	6.0	25%	4.2	92%	9.8	18%	4.4	87%	12.1	45%
2-Nonanol <sup>1</sup>	8	6.9	12%	5.3	22%	6.4	16%	32.7	87%	32.5	7%	35.9	8%
<b>Esters</b>													
2-Phenylethyl butanoate <sup>1,2</sup>	34	103.2	32%	20.6	26%	24.9	24%	167.5	10%	23.1	12%	23.2	12%
2-Phenethyl hexanoate <sup>1,2</sup>	35	1244.4	28%	74.5	22%	65.7	23%	1301.1	19%	70.6	22%	0.0	
Isoamyl propanoate <sup>1,2</sup>	19	10.8	10%	14.0	42%	20.2	14%	33.1	7%	29.4	1%	35.1	10%
Methyl 4-Methyl-2-hexenoate <sup>1,2</sup>	20	51.7	8%	48.7	14%	59.3	4%	128.6	1%	129.3	3%	153.2	1%
Methyl geranate <sup>1,2</sup>	21	35.2	23%	32.7	2%	39.4	2%	75.7	7%	77.2	12%	88.9	13%
isobutyl butanoate <sup>1</sup>	7	6.9	88%	6.1	87%	5.0	87%	22.8	7%	21.7	6%	25.4	4%
Ethyl hexanoate <sup>1,2</sup>	27	744.5	14%	571.9	11%	1999.8	18%	758.3	7%	548.2	9%	1249.8	10%
Ethyl octanoate <sup>1,2</sup>	28	797.7	18%	985.9	21%	1891.1	8%	438.6	14%	830.0	5%	1389.9	6%
Ethyl decanoate <sup>1,2</sup>	29	182.3	29%	195.1	28%	812.5	12%	46.3	15%	159.5	34%	703.1	18%
Ethyl dodecanoate <sup>1,2</sup>	30	26.3	35%	56.8	60%	129.2	15%	12.4	9%	58.8	66%	108.8	7%
<b>Higher Alcohols</b>													
2-Phenyl Ethanol <sup>1,2</sup>	38	1516.3	9%	862.4	10%	1103.9	18%	1962.5	17%	961.4	15%	1458.3	18%
Isoamyl alcohol <sup>2</sup>	10	751.3	6%	994.5	10%	1771.7	12%	525.0	87%	1070.4	5%	1796.5	4%
<b>Monoterpenes</b>													
2,7-Dimethyl-2,7-octadiene <sup>1</sup>	3	35.6	50%	29.8	55%	28.5	37%	56.0	7%	49.0	88%	18.0	94%
Citronellol <sup>1,2</sup>	14	35.5	18%	0.0		0.0		52.5	10%	41.4	19%	55.3	13%
Nerol <sup>1,2</sup>	13	0.0		0.0		0.0		8.0	24%	7.1	23%	10.2	20%
Linalool <sup>1</sup>	1	149.3	2%	146.1	2%	168.9	1%	342.2	6%	343.2	5%	352.3	5%
Geraniol <sup>1,2</sup>	12	12.0	3%	7.6	13%	10.1	18%	31.9	2%	19.2	2%	31.6	2%
<b>Sesquiterpenols</b>													
7-epi-amiteol <sup>1,2</sup>	17	105.2	6%	71.7	87%	0.0		108.8	13%	117.3	11%	28.7	173%
α/t-Cadinol <sup>1,2</sup>	18	0.0		17.0	11%	13.2	16%	0.0		84.0	13%	44.6	90%
Hop ether <sup>1</sup>	4	11.3	3%	11.6	17%	14.5	10%	23.2	9%	23.7	7%	23.5	10%
Juniper Camphor <sup>1</sup>	6	0.0		0.0		0.0		126.0	16%	129.3	12%	157.7	44%
β-Eudesmol <sup>1,2</sup>	15	0.0		0.0		0.0		163.8	14%	177.7	12%	151.2	15%
α-Eudesmol <sup>1,2</sup>	16	11.7	37%	11.6	26%	8.9	5%	168.5	15%	188.2	13%	161.1	16%
Caryophyllene oxide <sup>1</sup>	5	53.0	8%	47.3	7%	57.5	13%	63.1	15%	72.0	22%	73.8	27%
<b>Terpenyl acetates</b>													
Citronellol acetate <sup>1,2</sup>	24	8.6	87%	11.1	19%	19.0	8%	15.7	24%	23.9	26%	23.0	19%
Geranyl acetate <sup>1</sup>	2	16.1	12%	9.2	88%	15.5	11%	32.4	11%	34.2	12%	32.6	7%
Neryl acetate <sup>1,2</sup>	23	0.0		0.0		0.0		0.0		0.0		0.0	
<b>Phenols</b>													
4-Vinylguaiaicol <sup>1,2</sup>	25	31.3	39%	21.0	7%	573.2	7%	27.6	10%	18.9	20%	486.6	5%
Styrene <sup>2</sup>	11	2.2	173%	0.0		91.6	13%	6.4	41%	6.7	88%	78.7	18%
<b>Other</b>													
2,3-dihydro-benzofuran <sup>1,2</sup>	26	0.0		0.0		94.8	32%	0.0		0.0		90.1	98%

<sup>1</sup> compounds influenced by hop variety (p-value <0.05); <sup>2</sup> compounds influenced by yeast (p-value <0.05)



**Fig. 4** Graphical representation of the concentration of geraniol and linalool ( $\mu\text{g/L}$  rel 3-octanol) in beers produced with Aramis, Bouclier, Strisselspalt and Triskel as aromatic hop and fermented with yeast CLIB267, CLIB279 and UCD915

This observation confirms that isoamyl alcohol brought by hop in beer is negligible compared to those produced by yeast. In our condition, UCD915 reaches the highest concentration of isoamyl alcohol ( $\approx 1700 \mu\text{g/L}$  rel 3-octanol, Fig. 3).

According to Schwarz et al. [35], styrene (11) is a sweet-smelling fluid considered to be toxic for humans. In our conditions, beers produced by the yeast strain UCD915 contained high concentration of styrene (60 to 105  $\mu\text{g/L}$  rel 3-octanol) whereas beers produced by the two other yeast strains present a lower concentration for this compound (0 to 8  $\mu\text{g/L}$  rel 3-octanol). Styrene is produced by the same pathway than 4-vinylguaiacol (4VG) (25), by transformation of cinnamic and ferulic acids into styrene and 4VG respectively (Fig.1). Two processes for their production have been described in beer [36]: during wort boiling by thermal decarboxylation and during fermentation by enzymatic process if the yeast strain are "POF+" (Phenol Off-Flavor, Fig.1). The genes *PAD1* and *FDC1* of yeast *Saccharomyces cerevisiae* are involved in this process. Yeast strains that are able to decarboxylate cinnamic and ferulic acids are POF+. The styrene concentration that we observed strongly suggests that the UCD915 yeast strain is POF+ (Fig. 3) and would be in adequation with observations with Weissbier strains [14].

### 3.3 Impact of hop variety and yeast strain on beer volatile profile

Among 39 compounds measured in this experiment, 28 are influenced by both the hop varieties and yeast strains. Fifteen of these compounds are known to be hop-derived compounds, 3 monoterpenes, *ie* geraniol (12), nerol (13), citronellol (14); 4 sesquiterpenols *ie*  $\alpha$ -eudesmol (15),  $\beta$ -eudesmol (16), 7 epi amiteol (17), a/t cadinol (18), 3 esters, *ie* isoamylpropanoate (19), Methyl-4-methyl-2-hexenoate (20), methyl geranate (21), 2 alcohols, *ie* 2-undecanol (22), 2-hexanol (23), 2 terpenyl acetates, *ie* neryl acetate (24), citronellyl acetate (25) and one various compounds 2,3 dihydrobenzofuran (26). Thirteen of these compounds are yeast-derived compounds esters, *ie* ethyl hexanoate (27); ethyl octanoate (28); ethyl decanoate (29); ethyl dodecanoate (30); isoamylacetate (31); hexyl acetate (32); 2-phenethyl- acetate (33); 2-phenylethyl butanoate (34); 2-phenethyl hexanoate (35); three medium chain fatty acids : octanoic (36), decanoic (37) and dodecanoic acids (38), and 2-phenylethanol (39).

#### 3.3.1 Hop-derived compounds

##### 3.3.1.1 Monoterpenes

Geraniol (12) is an odour-active compound in beer [24] and is known to be hop variety dependent [11, 13, 19, 23, 24, 42, 43]. Yeast is able to transform geraniol into citronellol (14) [21, 41, 43] and to acetylate geraniol into geranyl acetate [41], but also other geraniol-derived compounds like nerol (13) and citronellol (14). We show here that hop and yeast together influenced the final composition of the beer. The maximum concentration which can be obtained is with the combination Aramis and CLIB267 (30  $\mu\text{g/L}$ ) (Fig.4).

##### 3.3.1.2 Sesquiterpenols

Sesquiterpenols concentration in beer is linked to the hop variety [40], however their influences on beer aroma are poorly understood.


We describe here that  $\alpha$ - and  $\beta$ -eudesmol (16-17), 7 epi amiteol (18), a/t cadinol (19) concentration (Table 1) are influenced by yeast. Beer fermented with ale yeast UCD915 is the less concentrated in sesquiterpenols, (Table 2 and Table 3). Our experiments also confirm that Aramis and Strisselspalt lead to higher sesquiterpenols production than Triskel or Bouclier [40] (Table 3 and Table 4).

##### 3.3.1.3 Esters


Isoamylpropanoate (19), methyl-4-methyl-2-hexenoate (20) and methyl geranate (21) have been reported in hop oil [2, 28, 29, 48] but their influences on beer aroma remain unclear. In our conditions there is a slight influence of yeast strain on both compounds while a strong influence of hop variety (Fig. 5). Surprisingly, isoamyl propanoate is not detected in beer produced with the yeast UCD915 using Aramis. Aramis can be considered as the highest producer of these three esters in beer (Table 1 and Fig. 5).

##### 3.3.1.4 Alcohols


As previously stated, 2-hexanol (22) and 2-undecanol (21) could be considered as the result of yeast-mediated 2-alkones, derived from iso- $\alpha$  acid from hop, reduction. It is not clear why and how 2-nonanol is influenced by yeast strain while 2-hexanol and 2-un-



**Fig. 5** Graphical representation of the concentration of methyl-4-methyl-2-hexenoate and isoamylpropanoate (µg/L rel 3-octanol) in beers produced with Aramis, Bouclier, Strisselspalt and Triskel as aromatic hop and fermented with yeast CLIB267, CLIB279 and UCD915



**Fig. 6** Graphical representation of the concentration of 4-vinylguaiacol (µg/L rel 3-octanol) and 2-phenylethanol in beers produced with Aramis, Bouclier, Strisselspalt and Triskel as aromatic hop and fermented with yeast CLIB267, CLIB279 and UCD915



**Fig. 7** Graphical representation of the concentration of ethyl hexanoate and ethyl octanoate (µg/L rel 3-octanol) in beers produced with Aramis, Bouclier, Strisselspalt and Triskel as aromatic hop and fermented with yeast CLIB267, CLIB279 and UCD915

decanol are (Table 4). To our knowledge there is no data on the influence of these compounds on the aromatic profile of beer. Both of these compounds are influenced by hop variety and yeast strain (Table 4).

2-phenylethanol (39), a higher alcohol and flavour active compound in beer, is known to be influenced by yeast and hop variety. Since it is an Erlich pathway by-product (Fig. 1), its concentration is linked to the yeast strain. Besides, it has been shown by Gros et

al. [26] that this compound can also be brought in beer by hop. In our case, Aramis and Triskel wort fermented with CLIB267 appears to be the combination bringing the highest concentration of 2-phenylethanol.

### 3.3.1.5 Various compounds

4 vinylguaiacol (4VG) (26) is a volatile compound with typical phenolic and clove aroma and is often found in Weissbier. This

compound is the result of the decarboxylation of ferulic acid by the enzymatic machinery of yeast (Fig.1). The yeast UCD915 is *POF+* and this hypothesis was confirmed by the high concentration of 4VG produced by this yeast in our conditions. Ferulic acid is a precursor of this compound and is provided by malt and hop. The malt used in our experiments remained unchanged in all 36 hop/yeast combinations. We conclude that the small variations of 4VG observed are linked to ferulic acid content of the hops. In our conditions, Triskel and Strisselspalt give the highest concentration of 4VG in beer (Fig.6).

### 3.3.2 Compounds deriving from yeast metabolism

Ethyl hexanoate (27); ethyl octanoate (28); ethyl dodecanoate (29); ethyl decanoate (30); isoamylacetate (31); hexyl acetate (32); 2-phenethylacetate (33); 2-phenylethylbutanoate (34); 2-phenethylhexanoate (35) are volatile compounds derived from yeast metabolism during fermentation. Most of them being flavour active, they could then contribute to the overall flavour of beer [31].

Surprisingly, production of these yeast derived compounds are influenced by the hop variety. Ethyl esters and acetate esters are significantly higher when Triskel hop was used (Fig.7). Neither of these esters (compounds 28–36) nor the precursors of these esters (1-hexanol, hexanoic, octanoic (36), decanoic (37), dodecanoic acids (38), 2-phenylethanol (39)) are known to be present in hop [5], or at best at very small concentrations (data not shown). However, it is well known that these compounds are produced by yeast during fermentation [34, 49] and are tightly linked to yeast genes involved in their production [20, 39]. As expected UCD915 (ale), produces high amount of acetate esters and ethyl esters compared to CLIB267 and 279 (lager yeasts) [33]. The same trends were also observed for medium chain fatty acids (compounds 36–38).

It has been shown that synthesis of esters can be repressed by presence of unsaturated fatty acids during fermentation [30, 32]. Interestingly, presence of unsaturated fatty acids have been reported in hop by *Demireva* et al. [8]. We propose that hop varieties contain different concentrations of these compounds which could influence in an indirect manner the production of esters by yeast. However there is a lack of data on the variability of these kind of compounds in hop. This hypothesis needs to be further investigated by measuring these unsaturated fatty acids in hops.

## 4 Conclusion

Hop varieties and yeast strains are often considered as key elements to distinguish beers. On one side, hop varieties are known to influence the concentration of various volatile compounds like monoterpenes and sesquiterpenols bringing floral and wood notes to the beer. On the other side, yeasts are known to be responsible for the productions of various esters giving fruity notes to the beer.

Some publications reported the influence of yeasts on specific volatile compounds brought by hop in beer [22] but none of them have evaluated the influence of different yeast strains combined with the use of different hop varieties on the volatile content of beer.

In this study, we evaluated the influence of yeast on the volatile compounds brought by hop but also to study the influence of hop on the volatile compounds produced by yeast.

We confirm here that some hop volatile compounds like linalool, geranylacetate, 2,7 dimethyl 2,7 octadiene, hop ether and juniper camphor are not metabolized by yeast during the fermentation. Nonetheless, some of them, including terpenols and sesquiterpenols, are influenced by yeast strains, for instance geraniol, neryl acetate, citronellol, b-eudesmol, 7 epi amiteol, a/t cadinol, isoamylpropanoate, methyl-4-methyl-2-hexanoate, 2,3 dihydrobenzofuran, 2-undecanol, 2-hexanol and 4 vinylguaiaicol.

Through different enzymes, yeasts are able to acetylate (*ATF1/ATF2*), reduce (*OYE2*) or decarboxylate (*PAD*) most of the volatiles or precursors of the volatiles found in this study. Mutation in enzyme gene(s) involved in this pathway is one way of explaining variability in yeast strains impact on volatile compounds. In other cases, like for 4VG, it could also depend on the concentration of the precursor provided, which is, in this case, hop-dependent.

Among all the volatile compounds produced by yeast, it has been shown in this study that compounds like ethyl esters, acetate esters and medium chain fatty acids are dependent on the yeast strain but also on the hop variety used. This interaction between hop variety and yeast strain could lead to better control the production of yeast-derived compounds in beer.

Hop can be used to monitor volatile compounds, for example, here we show that Aramis can be used to bring monoterpenol (floral) and sesquiterpenol (woody, spice), whereas Triskel and Strisselspalt increase the final concentration of 4-vinylguaiaicol (clove) when the appropriate yeast strain is used, finally we demonstrate here that Bouclier can bring neryl acetate (floral) in to the beer.

We propose here that a separate view of yeast and hop input for beer flavouring might be too restrictive. The intricacy of yeast/hop combination allows for a more complex and subtle control of the beer produced and should be assessed for an advanced understanding of beer design.

## Acknowledgments

We thank *Patricia Claudel* (INRA Colmar) for the GC/MS analysis, *Anne Alais* (INRA Colmar) for the kindly provision of yeast strain collection, *Jean-Luc Legras* (INRA Montpellier), *Ann-Marie Mc Cartin* (University College Dublin), *Valentin Le Douce* (Emosis, Illkirch-Graffenstaden) for their support of this work.

## 5 Literature

1. Bekatorou, A.; Sarellas, A.; Ternan, N. G.; Mallouchos, A.; Komaitis, M. and Koutinas, A.: Low-Temperature Brewing Using Yeast Immobilized on Dried Figs, *J. Agric. Food Chem.*, **50** (2002), pp. 7249-7257.
2. Briggs, D. E.; Brookes, P.; Stevens, R. and Boulton, C.: *Brewing: Science and Practice*, (Woodhead Publishing, 2004).
3. Burdock, G. A.: Fenaroli's Handbook of Flavor Ingredients, Fourth Edition, in Fenaroli's Handb. Flavor Ingredients, Fourth Ed., (CRC

- Press, 2001), p. 147.
4. Carrau, F.; Boido, E. and Dellacassa, E.: Yeast Diversity and Flavor Compounds, in *Fungal Metab.*, (2015), pp. 1-29.
  5. Cibaka, M.-L. K.; Gros, J. and Collin, S.: Revue sur les étonnantes analogies et les différences relevées entre un cône de houblon et une baie de raisin, *Cerevisia*, **38** (2014), pp. 103-117.
  6. Coelho, E.; Coimbra, M.; Nogueira, J. M. F. and Rocha, S. M.: Quantification approach for assessment of sparkling wine volatiles from different soils, ripening stages, and varieties by stir bar sorptive extraction with liquid desorption., *Anal. Chim. Acta*, **635** (2009), pp. 214-21.
  7. Daenen, L.; Sterckx, F.; Delvaux, F. R.; Verachert, H. and Derdelinckx, G.: Evaluation of the glycoside hydrolase activity of a *Brettanomyces* strain on glycosides from sour cherry (*Prunus cerasus* L.) used in the production of special fruit beers., *FEMS Yeast Res.*, **8** (2008), pp. 1103-14.
  8. Demireva, Z. N. and Totova, P. F. : Studies on the Fatty Acid Composition of Hop Extract, *J. Inst. Brew.*, **101** (1995), pp. 437-438.
  9. Tressl, R.; Engel K. H.; Kossa, M. and H. Koeppler, H: Characterization of tricyclic sesquiterpenes in hop (*Humulus lupulus* var. Hersbrucker Spaet), *J. Agric. Food Chem.*, **31** (1983), no. 4, pp. 4758-64.
  10. Eri, S.; Khoo, B. K., Lech, J. and Hartman, T. G.: Direct Thermal Desorption–Gas Chromatography and Gas Chromatography–Mass Spectrometry Profiling of Hop (*Humulus lupulus* L.) Essential Oils in Support of Varietal Characterization, *J. Agric. Food Chem.*, **48** (2000), pp. 1140-1149.
  11. Van Opstaele, F.; De Rouck, G.; De Clippeleer, J.; Aerts, G. and De Cooman, L.: Analytical and Sensory Assessment of Hoppy Aroma and Bitterness of Conventionally Hopped and Advanced Hopped Pilsner Beers, *J. Inst. Brew.*, **116** (2010), no. 4, pp. 445-458.
  12. Forster, A.; Gahr, A. and Van Opstaele, F.: On the Fate of Certain Hop Substances during Dry Hopping, **66** (2013), pp. 93-103.
  13. Forster, A.; Gahr, A. and Van Opstaele, F.: On the Transfer Rate of Geraniol with Dry When calculating the transfer rate of geraniol in dry hopped beers based solely on the initial value in hops, **67** (2014), pp. 60-62.
  14. Gallone, B.; Steensels, J.; Prahl, T.; Soriaga, L.; Saels, V. and Herrera-Malaver, B.: Domestication and Divergence of *Saccharomyces cerevisiae* Beer Yeasts, *Cell*, **166** (2016), pp. 1397-1410.
  15. Goiris, K.; Ridder, M.; Rouck, G.; Boeykens, A.; Opstaele, F.; Aerts, G.; De Cooman, L. and De Keukeleire, D.: The Oxygenated Sesquiterpenoid Fraction of Hops in Relation to the Spicy Hop Character of Beer, *J. Inst. Brew.*, **108** (2002), pp. 86-93.
  16. Hanke, S.: Linalool – A Key Contributor to Hop Aroma, *Global Emerging Issues* (2009).
  17. Hashimoto, N.: Contribution of hop bitter substances to head formation of beer.pdf, *J. Am. Soc. Brew. Chem.*, **37** (1979).
  18. Hashimoto, N. and Kuroiwa, Y.: Proposed Pathways for the Formation of Volatile Aldehydes during Storage of Bottled Beer, *Am. Soc. Brew. Chem. Proc.*, (1975), pp. 104-111.
  19. Inui, T.; Tsuchiya, F.; Ishimaru, M.; Oka, K. and Komura, H.: Different beers with different hops. Relevant compounds for their aroma characteristics., *J. Agric. Food Chem.*, **61** (2013), pp. 4758-64.
  20. Katou, T.; Namise, M.; Kitagaki, H.; Akao, T. and Shimoi, H.: QTL mapping of sake brewing characteristics of yeast, *J. Biosci. Bioeng.*, **107** (2009), pp. 383-393.
  21. King, A. and Dickinson, J. R.: Biotransformation of monoterpene alcohols by *Saccharomyces cerevisiae*, *Torulaspota delbrueckii* and *Kluyveromyces lactis*., *Yeast*, **16** (2000), pp. 499-506.
  22. King, A. J. and Dickinson, J. R.: Biotransformation of hop aroma terpenoids by ale and lager yeasts, *FEMS Yeast Res.*, **3** (2003), pp. 53-62.
  23. Kishimoto, T.; Wanikawa, A.; Kagami, N. and Kawatsura, K.: Analysis of hop-derived terpenoids in beer and evaluation of their behavior using the stir bar-sorptive extraction method with GC-MS., *J. Agric. Food Chem.*, **53** (2005), pp. 4701-4707.
  24. Kishimoto, T.; Wanikawa, A.; Kono, K. and Shibata, K.: Comparison of the Odor-Active Compounds in Unhopped Beer and Beers Hopped with Different Hop Varieties, *J. Agric. Food Chem.*, **54** (2006), pp. 8855-8861.
  25. Lam, K. and Deinzer, M.: Contribution of hop ether and Karahana ether to beer flavor, *J. Am. Soc. Brew. Chem.*, **44** (1986), pp. 69-72.
  26. Lermusieau, G.; Bulens, M. and Collin, S.: Use of GC – Olfactometry to Identify the Hop Aromatic Compounds in Beer, *J. Agric. Food Chem.*, **49** (2001), no. 8, pp. 3867-3874
  27. Lermusieau, G. and Collin, S.: Hop Aroma Extraction and Analysis, *Anal. Tast. Aroma*, (2002), pp. 69-88.
  28. McAdam, E. L.; Freeman, J. S.; Whittock, S. P.; Buck, E. J.; Jakse, J.; Cerenak, A. et al.: Quantitative trait loci in hop (*Humulus lupulus* L.) reveal complex genetic architecture underlying variation in sex, yield and cone chemistry., *BMC Genomics*, **14** (2013), pp. 360.
  29. Nance; M. R. and Setzer, W. N.: Volatile components of aroma hops (*Humulus lupulus* L.) commonly used in beer brewing, *Journal of brewing and distilling*, **2** (2011), pp. 16-22.
  30. Peddie, H.: Ester formation in brewery fermentations, *J. Inst. Brew.*, **96** (1990), pp. 327-331.
  31. Preedy, V. R. V.: Beer in Health and Disease Prevention, (2009).
  32. Saerens, S. M. G.; Delvaux, F.; Verstrepen, K. J.; Dijck, P. Van, Thevelein; J. M. and Delvaux, F. R.: Parameters affecting ethyl ester production by *Saccharomyces cerevisiae* during fermentation., *Appl. Env. Microbiol.*, **74** (2008), pp. 454-461.
  33. Saerens, S. M. G.; Verbelen, P. J.; Vanbeneden, N.; Thevelein, J. M. and Delvaux, F. R.: Monitoring the influence of high-gravity brewing and fermentation temperature on flavour formation by analysis of gene expression levels in brewing yeast, *Appl. Microbiol. Biotechnol.*, **80** (2008), pp. 1039-1051.
  34. Saerens, S. M. G.; Verstrepen, K. J.; Van Laere, S. D. M.; Voet, A. R. D.; Van Dijck, P.; Delvaux, F. R.; et al.: The *Saccharomyces cerevisiae* EHT1 and EEB1 genes encode novel enzymes with medium-chain fatty acid ethyl ester synthesis and hydrolysis capacity., *J. Biol. Chem.*, **281** (2006), pp. 4446-56.
  35. Schwarz, K.: Styrene Concentrations during Wheat Beer Production, **64** (2011), pp. 156-160.
  36. Schwarz, K. J.; Stübner, R. and Methner, F. J.: Formation of styrene dependent on fermentation management during wheat beer production, *Food Chem.*, **134** (2012), pp. 2121-2125.
  37. Sharpe, B. F. R.; Laws, D. R. J.; Brewer; N.; Sharpe; F. and Laws, D. R. J.: The essential oil of hops a review, *J. Inst. Brew.*, **87** (1981) , pp. 96-107.
  38. Sharp, D.C.: Harvest Maturity of Cascade and Willamette Hops, (2013).
  39. Steyer, D.; Ambroset, C.; Brion, C.; Claudel, P.; Delobel, P.; Sanchez, I.; et al.: QTL mapping of the production of wine aroma compounds by yeast., *BMC Genomics*, **13** (2012) , pp. 573.
  40. Steyer, D.; Clayeux, C. and Laugel, B.: Characterization of the terpenoids composition of beers made with the French hop varieties: Strisselspalt, Aramis, Triskel and Bouclier, *BrewingScience*, **66** (2013), no. 11/12, pp. 192-197.
  41. Steyer, D.; Erny, C.; Claudel, P.; Riveill, G.; Karst, F. and Legras, J.: Genetic analysis of geraniol metabolism during fermentation., *Food Microbiol.*, **33** (2013), pp. 228-34.

42. Takoi, K.; Itoga, Y.; Koie, K.; Kosugi, T.; Katayama, Y.; Nakayama, Y.; et al.: The Contribution of Geraniol Metabolism to the Citrus Flavour of Beer : Synergy of Geraniol and  $\beta$ -Citronellol Under Coexistence with Excess Linalool, *Journal of the Institute of Brewing*, **116** (2010), pp. 3-5.
43. Takoi, K.; Koie, K.; Itoga, Y.; Katayama, Y.; Shimase, M.; Nakayama, Y.; et al.: Biotransformation of hop-derived monoterpene alcohols by lager yeast and their contribution to the flavor of hopped beer., *J. Agric. Food Chem.*, **58** (2010), pp. 5050-8.
44. Tressl, R.; Friese, L.; Fendesack, F. and Koppler, H.: Studies of the Volatile Composition of Hops during Storage, *26* (1978), pp. 1426-1430.
45. Tressl, R., Friese, L., Fendesack, F., and Koppler, H., GC/MS Investigation of hop aroma constituents in beer., *J. Agric. Food Chem.*, **26** (1978), pp. 1422-1426.
46. Tsuji, H. and Mizuno, A., Volatile Compounds and the Changes in Their Concentration Levels during Storage in Beers Containing Varying Malt Concentrations, *J. Food Sci.*, **75** (2010), pp. 79-84.
47. Vanderhaegen, B.; Neven, H.; Verachtert, H. and Derdelinckx, G.: The chemistry of beer aging – a critical review, *Food Chem.*, **95** (2006), pp. 357-381.
48. Vázquez-Araújo, L.; Rodríguez-Solana, R.; Cortés-Diéguez, S. M. and Domínguez, J. M.: Use of hydrodistillation and headspace solid-phase microextraction to characterize the volatile composition of different hop cultivars., *J. Sci. Food Agric.*, **93** (2013), pp. 2568-74.
49. Verstrepen, K. J.; Derdelinckx, G.; Dufour, J. P.; Winderickx, J.; Pretorius, I. S.; Thevelein, J. M.; et al.: The *Saccharomyces cerevisiae* alcohol acetyl transferase gene ATF1 is a target of the cAMP/PKA and FGM nutrient-signalling pathways., *FEMS Yeast Res.*, **4** (2003), pp. 285-296.

Received 16 August 2017, accepted 16 October 2017